

Implicit-explicit methods based on strong stability preserving multistep time discretizations[★]

Thor Gjesdal

Norwegian Defence Research Establishment, NO-2027 Kjeller, Norway

Abstract

In this note we propose and analyze novel implicit-explicit methods based on second order strong stability preserving multistep time discretizations. Several schemes are developed, and a linear stability analysis is performed to study their properties with respect to the implicit and explicit eigenvalues. One of the proposed schemes is found to have very good stability properties, with implicit A -stability for the entire explicit stability domain. The properties of the other proposed schemes are comparable to those of traditional methods found in the literature.

Key words: Implicit-explicit methods, strong stability preserving methods, advection-diffusion equation, stability

1 Introduction

Implicit-explicit schemes (IMEX) are methods for the solution of time-dependent differential equations in the form

$$\dot{y} = f(y, t) + g(y, t), \quad (1)$$

where f and g are terms with different character such that one, say g , mandates implicit treatment whereas $\dot{y} = f$ can be solved efficiently by an explicit method. Such systems often arise from spatial (semi-)discretization of time

[★] This work was initiated while the author was employed by Christian Michelsen Research AS (CMR), Bergen, Norway; and supported in part by the sponsors of CMR's Gas Safety Programme (GSP 97–99): Agip, British Petroleum, Elf, Exxon, Gaz de France, Health and Safety Executive, Mobil, Norsk Hydro, Norwegian Petroleum Directorate, Phillips Petroleum, and Statoil.

dependent PDEs by the method-of-lines – a prime example being discretizations of transport equations that may contain different terms accounting for advection, diffusion, and reaction.

It is beyond the scope of this note to give a complete description of IMEX methods; for a systematic discussion of multistep IMEX schemes we refer to Ascher et al. [2]. Frank et al. [3] analyzed the stability of multistep IMEX schemes, and showed that stable implicit and explicit integrators do not necessarily lead to a stable implicit-explicit method. Ascher et al. [1] have also developed IMEX schemes based on multistage Runge-Kutta integrators.

The Strong Stability Preserving (SSP) property is a generalization of the Total Variation Diminishing (TVD) property for hyperbolic problems, and it is a guarantee of non-linear stability of the scheme. Traditional explicit multistep schemes, such as the Adams-Bashforth methods, do not necessarily have the SSP property [4]. Note however that recent work – by Hundsdorfer and Jaffré [6] and by Hundsdorfer, Ruuth, and Spiteri [8] – have shown that the traditional multistep methods can be made monotonicity-preserving, e.g. with respect to the Total Variation semi-norm, if a suitable starting procedure is employed.

In this note we discuss the linear stability of novel IMEX schemes based on second order accurate SSP multistep time discretization. Although SSP is a non-linear stability property, it is nevertheless appropriate to study the linear stability of the proposed schemes, since IMEX combinations do not necessarily inherit the stability properties of the component schemes [3]. A full analysis of the nonlinear stabilities of the proposed schemes is beyond the scope of this note.

We will show that second order IMEX schemes based on the explicit multistep SSP discretizations proposed by Shu [11] have stability properties that are comparable to the Crank-Nicholson/Adams-Bashforth and BDF2 IMEX methods.

2 Stability analysis

A general implicit-explicit k -step method for (1) can be written in the form

$$\sum_{i=0}^k a_i y_{n+1-i} = \sum_{i=1}^k b_i f_{n+1-i} + \sum_{i=0}^k c_i g_{n+1-i}, \quad (2)$$

where $f_n = f(y_n, t_n)$ and $g_n = g(y_n, t_n)$.

The stability analysis below is based on the scalar test equation

$$\dot{y} = \lambda y + \mu y, \quad (3)$$

where λ and μ are complex constants that represent the eigenvalues of the explicit and implicit operators, respectively.

By applying the multistep method (2) to the scalar test equation (3) and looking for solutions in the form $y = \zeta^n$, we obtain the characteristic equation

$$\sum_{i=0}^k a_i \zeta^{n+1-i} - \lambda \sum_{i=1}^k b_i \zeta^{n+1-i} - \mu \sum_{i=0}^k c_i \zeta^{n+1-i} = 0. \quad (4)$$

For convenience, we follow Frank et al. [3] in working on the transformed form of the equation

$$A(z) - \lambda B(z) - \mu C(z) = 0, \quad (5)$$

where $z = 1/\zeta$ and the polynomials A , B , and C correspond to the time derivative, explicit, and implicit operators, respectively, and are given by

$$A(z) = \sum_{i=0}^k a_i z^i, \quad B(z) = \sum_{i=0}^k b_i z^i, \quad C(z) = \sum_{i=0}^k c_i z^i. \quad (6)$$

Note that, since we are working in the transformed variable z , the scheme (2) is stable if all the roots of the characteristic equation (5) are in the exterior of the unit disk, $|z| \geq 1$ (with strict inequality if z is a multiple root).

To investigate the stability properties of the combined IMEX scheme we will study the image of the exterior of the unit circle under the mapping

$$\mu = \varphi_\lambda(z) = \frac{A(z) - \lambda B(z)}{C(z)}. \quad (7)$$

The boundary of the stability domain, $\mathcal{S}$, for the explicit method can be readily determined from

$$\partial\mathcal{S} = \left\{ \lambda : \lambda(\theta) = A(e^{i\theta})/B(e^{i\theta}), \quad -\pi \leq \theta < \pi \right\}.$$

Similarly, we can find the stability region, $\mathcal{D}$, of the implicit method from

$$\partial\mathcal{D} = \left\{ \mu : \mu(\theta) = A(e^{i\theta})/C(e^{i\theta}), \quad -\pi \leq \theta < \pi \right\}.$$

3 IMEX methods based on second order multistep SSP schemes

3.1 SSP time discretizations

The first order accurate forward Euler explicit time integration is strongly stable for some norm, $\|\cdot\|$, if

$$\|y_n + \Delta t f_n\| \leq \|y_n\|,$$

under a suitable time step $\Delta t \leq \Delta t_0$, where Δt_0 is the largest allowable time step for stability. Higher order time discretizations are strong stability preserving (SSP) if they retain this property, possibly for a shorter time step $\Delta t \leq c\Delta t_0$. The constant $c \leq 1$ is, by convention, called the CFL coefficient [4], and should not be confused with the Courant number of hyperbolic discretizations. The SSP time discretizations were originally developed by Shu [11] for the Total Variation norm to maintain positivity and monotonicity in the solution of hyperbolic equations.

Shu [11] proposed a family of explicit SSP multistep schemes. We will restrict the discussion to second order schemes with positive coefficients. We thus consider the three- and four-step second order schemes:

$$\dot{y}_{\Delta t}^{[3]} = \frac{4y_{n+1} - 3y_n - y_{n-2}}{6\Delta t} = f_n, \quad (8a)$$

$$\dot{y}_{\Delta t}^{[4]} = \frac{9y_{n+1} - 8y_n - y_{n-3}}{12\Delta t} = f_n, \quad (8b)$$

where $\dot{y}_{\Delta t}^{[k]}$ denotes the k -step discretized time derivative for the SSP scheme. These methods are strong stability preserving with CFL coefficients $c = 1/2$ and $c = 2/3$ for $k = 3$ and $k = 4$, respectively. The polynomials (6) that characterize these schemes are

$$A^{[3]}(z) = (4 - 3z - z^3)/6, \quad (9a)$$

$$A^{[4]}(z) = (9 - 8z - z^4)/12, \quad (9b)$$

and

$$B(z) = z. \quad (10)$$

We show the explicit stability domain, $\mathcal{S}$, for the two schemes (8) in Fig. 1. Notice that both schemes have an appreciable part of the stability region close to the imaginary axis, something that is advantageous for the solution of hyperbolic terms.




Fig. 1. Explicit stability domain, $\mathcal{S}$, for the two SSP time discretizations (8).

3.2 Construction of implicit integrators

To develop IMEX methods based on the multistep SSP schemes, we first construct implicit schemes by replacing the term $f(y^n)$ in (8) above by linear combinations of the implicit operator evaluated at the available time levels

$$\sum_{j=0}^k c_j g_{n+1-j},$$

such that the resulting formula for calculating y^{n+1} is second order accurate at time level n . We must obviously require $c_0 \neq 0$ for the scheme to be implicit, and furthermore that $\sum c_j = 1$.

Evaluating the order conditions we obtain an under-determined system that allows infinitely many solutions with several free parameters. We will not go into all the possibilities here. Instead we focus on the straightforward approximations

$$\dot{y}_{\Delta t}^{[k]} = \frac{1}{3} \left(2g_{n+1} + g_{n-2} \right), \quad (11a)$$

$$\dot{y}_{\Delta t}^{[k]} = \frac{1}{2} \left((1 - \beta)g_{n+1} + 2\beta g_n + (1 - \beta)g_{n-1} \right), \quad (11b)$$

where $\beta < 1$ is a non-negative algorithmic parameter.

The polynomials (6) that correspond to the right-hand-sides in (11) are then

$$C_1(z) = (2 + z^3)/3, \quad (12a)$$

$$C_2(z) = \left((1 - \beta) + 2\beta z + (1 - \beta)z^2 \right) / 2. \quad (12b)$$

An implicit method has A -stability if it is stable for all eigenvalues, μ , in the right half plane, $\text{Re}(\mu) \leq 0$, and it has $A(\alpha)$ -stability if it is stable for eigenvalues in a wedge-shaped domain about the negative real axis with half-angle α .

In Fig. 2 we show the stability regions of the $k = 3$ versions of the implicit integrators (11). Note that the scheme (11a) appears to be A -stable, whereas the other variant (11b) is $A(\alpha)$ -stable. We can show that these stability properties carry over to the $k = 4$ versions of the schemes. That is, (11a) is A -stable and (11b) is $A(\alpha)$ -stable, for both $k = 3$ and $k = 4$.

Lemma 1 *The implicit method (11a) is A -stable for both $k = 3$ and $k = 4$.*

PROOF. Obviously, for a purely implicit method $\lambda = 0$, and we consider the mapping

$$\varphi_0(z) = A^{[k]}(z)/C_1(z).$$

A -stability requires that the image of the unit circle under this mapping is entirely in the right half plane, i.e. $\text{Re}(\varphi_0(e^{i\theta})) \geq 0$.

(i) For $k = 3$ we find

$$\text{Re}(\varphi_0(e^{i\theta})) = \frac{14 - 12 \cos \theta - 6 \cos 2\theta + 4 \cos 3\theta}{4 |2 + e^{i3\theta}|^2} \quad (13)$$

$$= \frac{5 - 6 \cos \theta - 3 \cos^2 \theta + 4 \cos^3 \theta}{|2 + e^{i3\theta}|^2}. \quad (14)$$

The denominator in this expression is obviously positive, we must therefore show that the numerator is non-negative. Let $x = \cos \theta$ and write

$$f(x) = 4x^3 - 3x^2 - 6x + 5 = (4x - 5)(x - 1)^2.$$

It is straightforward to verify that $f(x)$ is non-negative for $-1 \leq x \leq 1$, and we have shown that (11a) is A -stable for $k = 3$.

(ii) Similarly, for $k = 4$, we have

$$\varphi_0(e^{i\theta}) = A^{[4]}(e^{i\theta})/C_1(e^{i\theta}),$$

leading to

$$\operatorname{Re}(\varphi_0(e^{i\theta})) = \frac{6 - 11 \cos \theta + 9 \cos^3 \theta - 4 \cos^4 \theta}{4 |2 + e^{i3\theta}|^2}.$$

Again we let $x = \cos \theta$ and write the numerator as

$$f(x) = -4x^4 - 9x^3 - x + 6 = -(4x^2 - x - 6)(x - 1)^2,$$

which we readily verify is non-negative for $-1 \leq x \leq 1$. $\square$

Lemma 2 *The implicit method (11b) is $A(\alpha)$ -stable, with $\alpha = \alpha_0$ given by*

$$\tan \alpha_0 = \frac{(2 + \gamma)\sqrt{1 - \gamma^2}}{(\gamma - 1)^2}, \quad \text{for } k = 3, \quad (15a)$$

$$\tan \alpha_0 = \frac{(2 + \gamma^2)\sqrt{1 - \gamma^2}}{2 - 3\gamma + \gamma^3}, \quad \text{for } k = 4, \quad (15b)$$

where $\gamma = \beta/(\beta - 1)$.

PROOF. Again, we let $\lambda = 0$, and we consider the mapping

$$\varphi_0(e^{i\theta}) = A^{[k]}(e^{i\theta})/C_2(e^{i\theta}),$$

shown in Fig. 2, to determine the half-angle, α_0 . Note that $\varphi_0(\exp(i\theta))$ is singular in the limit $\cos \theta \rightarrow \gamma = \beta/(\beta - 1)$. We only consider $\beta \leq 1/2$; since for $\beta > 1/2$ the scheme behaves more like an explicit method. The angle α_0 is given by the slope of the asymptote, which can be written

$$\tan \alpha_0 = \lim_{\cos \theta \rightarrow \gamma} \frac{\operatorname{Im}(\varphi_0(e^{i\theta}))}{\operatorname{Re}(\varphi_0(e^{i\theta}))},$$

We then have

$$\begin{aligned} \tan \alpha_0 &= \frac{-4 \sin \theta^* - \sin 3\theta^*}{4 - 3 \cos \theta^* - \cos 3\theta^*}, & \text{for } k = 3, \\ \tan \alpha_0 &= \frac{-8 \sin \theta^* - \sin 4\theta^*}{9 - 8 \cos \theta^* - \cos 4\theta^*}, & \text{for } k = 4, \end{aligned}$$

from which the results follow. $\square$

The natural choice $\beta = 0$, i.e. $\gamma = 0$, gives $\tan \alpha_0 = 2$ and $\tan \alpha_0 = 1$ for the three-step and the four-step schemes respectively. In both cases, this is the optimal parameter value that maximizes α_0 .




Fig. 2. Stability regions, $\mathcal{D}$, for the implicit integrators (11), where $k = 3$ and $\beta = 0$. The methods are stable outside the shaded regions.

3.3 Stability of the IMEX schemes

We can construct implicit explicit methods by combining the above SSP explicit schemes (8) and the implicit integrators (11). Thus we consider

$$\dot{y}_{\Delta t}^{[k]} = f(y^n) + \frac{1}{3} \left(2g(y^{n+1}) + g(y^{n-2}) \right). \quad (16a)$$

$$\dot{y}_{\Delta t}^{[k]} = f(y^n) + \frac{1}{2} \left((1 - \beta)g(y^{n+1}) + 2\beta g(y^n) + (1 - \beta)g(y^{n-1}) \right), \quad (16b)$$

Frank et al. [3] presented two general stability results for multistep IMEX schemes. The first gives restrictions on the explicit eigenvalues in order to retain the full stability domain of the implicit operator, whereas the second gives restrictions on the implicit eigenvalues in order to retain the full stability domain of the explicit operator. Unfortunately, neither of these results appear to be applicable to the proposed schemes, and we will therefore study the stability properties of (16) by working directly on the mapping $\varphi_\lambda(\exp(i\theta))$ from (7). Remember that the IMEX schemes are stable with respect to implicit eigenvalues, μ , that are in the exterior of the mapping of the unit disk under φ_λ . For a method to be A -stable, the entire image of the unit circle must be in the right half-plane.

3.3.1 Stability of scheme (16a)

In this section we will discuss the stability of the scheme (16a), characterized by the polynomials (9), (10), and (12a). For the $k = 3$ variant of the scheme we propose the following stability result:

Conjecture 3 *The scheme (16a) with $k = 3$ is A -stable with respect to μ for $\lambda \in \mathcal{S}$.*

Recall that A -stability requires that the image of the unit circle under the mapping $\varphi_\lambda(z)$ must be entirely in the right half-plane for $\lambda \in \mathcal{S}$, i.e.

$$\Phi(\lambda, z) = \operatorname{Re}(\varphi_\lambda(e^{i\theta})) \geq 0, \quad -\pi \leq \theta < \pi, \quad \lambda \in \mathcal{S}.$$

As a consequence of the maximum principle, Φ will attain its minimal value w.r.t. λ on the boundary of the explicit stability domain. We need therefore only consider $\lambda \in \partial\mathcal{S}$.

To support the assertion that (16a) with $k = 3$ is A -stable, we first show, in Fig. 3, the image of the unit circle under φ_λ for $\lambda \in \partial\mathcal{S}$. Furthermore, let $\lambda^* = A^{[3]}(\exp(i\theta^*))/B(\exp(i\theta^*))$ be an eigenvalue on the boundary of the explicit stability domain. Then $\varphi_{\lambda^*}(\exp(i\theta^*)) = 0$, and we can investigate the behaviour of φ_{λ^*} close to this zero by performing a Taylor expansion. To leading order we obtain

$$\operatorname{Re}(\varphi_{\lambda^*}(e^{i\theta})) \approx \frac{1 - \cos 3\theta^*}{5 + 4 \cos 3\theta^*}(\theta - \theta^*)^2 \geq 0. \quad (17)$$

We have neither analytically nor numerically managed to find other zeros or negative values for $\Phi(\lambda, z)$ and the scheme does therefore indeed appear to be A -stable.

For the $k = 4$ variant we propose:

Conjecture 4 *The scheme (16a) with $k = 4$ is $A(\alpha)$ -stable with respect to μ for $\lambda \in \mathcal{S}$. The half-angle, α , of the wedge of stability is $\alpha \approx 0.23\pi$.*

In Fig. 4 we show the image of the unit circle under φ_λ for $\lambda \in \partial\mathcal{S}$, and we observe that the method version appears to be $A(\alpha)$ stable with $\alpha \approx \pi/4$. Carrying out a Taylor expansion of φ_{λ^*} as above, we find to leading order

$$\operatorname{Re}(\varphi_\lambda(e^{i\theta})) \approx \frac{3}{4} \frac{\sin \theta^* - 3 \sin 3\theta^* + 2 \sin 4\theta^*}{\sin^2 3\theta^* + (\cos 3\theta^* + 2)^2}(\theta - \theta^*), \quad (18a)$$

$$\operatorname{Im}(\varphi_\lambda(e^{i\theta})) \approx \frac{3}{4} \frac{6 + \cos \theta^* + 3 \cos 3\theta^* + 2 \cos 4\theta^*}{\sin^2 3\theta^* + (\cos 3\theta^* + 2)^2}(\theta - \theta^*). \quad (18b)$$




Fig. 3. Image of the unit circle under the mapping $\varphi_\lambda(z)$, with $\lambda \in \partial\mathcal{S}$, for the schemes given by (16a) with $k = 3$. The implicit stability domain of the method is outside the shaded region.

We can estimate the angle for $A(\alpha)$ -stability from the slope of the first order approximation:

$$\tan \alpha \approx \inf_{\theta^*} \left| \frac{6 + \cos \theta^* + 3 \cos 3\theta^* + 2 \cos 4\theta^*}{\sin \theta^* - 3 \sin 3\theta^* + 2 \sin 4\theta^*} \right|. \quad (19)$$

Numerically we find that $\tan \alpha \approx 0.89$, and this gives $\alpha \approx 0.23\pi$ which corresponds well with the stability domain shown in Fig. 4.

3.3.2 Stability of scheme (16b)

We then consider the scheme (16b), characterized by the polynomials (9), (10), and (12b). For this scheme, we can show the following stability result:

Lemma 5 *The implicit-explicit method (16b) is $A(\alpha)$ -stable with respect to the implicit eigenvalues, μ , for $\beta \leq \frac{1}{2}$ if*

$$\lambda \in \mathcal{S}^\nu = \mathcal{S} \cap \{z : |\operatorname{Im}(z)| \leq \nu\}, \quad \text{and} \quad |\alpha| \leq \alpha_\nu$$




Fig. 4. Image of the unit circle under the mapping $\varphi_\lambda(z)$, with $\lambda \in \partial\mathcal{S}$, for the schemes given by (16a) with $k = 4$. The implicit stability domain of the method is outside the shaded region.

where ν and α_ν are given by

$$\nu \leq (2 + \gamma)\sqrt{1 - \gamma^2}/3, \quad (20a)$$

$$\tan \alpha_\nu = \frac{(2 + \gamma)\sqrt{1 - \gamma^2} - 3\nu}{(\gamma - 1)^2}, \quad (20b)$$

for $k = 3$, and

$$\nu \leq (2 + \gamma^2)\sqrt{1 - \gamma^2}/3, \quad (21a)$$

$$\tan \alpha_\nu = \frac{(2 + \gamma^2)\sqrt{1 - \gamma^2} - 3\nu}{2 - 3\gamma + \gamma^3}. \quad (21b)$$

for $k = 4$.

PROOF. Recall from Sec. 3.2 that, in this case, the mapping $\varphi_\lambda(\exp(i\theta))$ is singular in the limit $\cos \theta \rightarrow \gamma = \beta/(\beta - 1)$. Taking into account a non-zero $\lambda = \lambda_r + i\lambda_i \in \mathcal{S}$ and proceeding as in the analysis of Sec. 3.2, we have

$$\tan \alpha_\lambda = \inf_{\lambda} \left(\frac{(2 + \gamma)\sqrt{1 - \gamma^2} - 3\lambda_i}{(\gamma - 1)^2 + 3\lambda_r} \right), \quad \text{for } k = 3$$

$$\tan \alpha_\lambda = \inf_{\lambda} \left(\frac{(2 + \gamma^2)\sqrt{1 - \gamma^2} - 3\lambda_i}{\gamma^3 - 3\gamma + 2 + 3\lambda_r} \right) \quad \text{for } k = 4.$$

We observe that for large $|\lambda_i|$, say $|\lambda_i| > \nu$; $\tan \alpha_\lambda$ may become negative. This means that the wedge of stability will be rotated entirely into the upper or lower left quadrants. In order to have a useful stability domain that – at least – contains the negative real axis, $\tan \alpha_\lambda$ must obviously be non-negative, and we must introduce an upper limit on the imaginary part of the explicit eigenvalues,

$$\begin{aligned} |\lambda_i| &\leq \nu \leq (2 + \gamma)\sqrt{1 - \gamma^2}/3, & \text{for } k = 3 \\ |\lambda_i| &\leq \nu \leq (2 + \gamma^2)\sqrt{1 - \gamma^2}/3, & \text{for } k = 4. \end{aligned}$$

We obtain a lower bound on the angle α if we choose $\lambda = \pm i\nu$, and we have:

$$\begin{aligned} \tan \alpha_\nu &= \frac{(2 + \gamma)\sqrt{1 - \gamma^2} - 3\nu}{(\gamma - 1)^2}, \\ \tan \alpha_\nu &= \frac{(2 + \gamma^2)\sqrt{1 - \gamma^2} - 3\nu}{2 - 3\gamma + \gamma^3}. \end{aligned}$$

for $k = 3$ and $k = 4$, respectively. $\square$

Note that the natural choice $\beta = 0$ maximizes both the angle α and the upper bound on ν , and thus gives the largest stability domains for both the implicit and explicit operators. Furthermore, although the above results require that we use a reduced explicit stability domain, the restrictions are not prohibitive. The choice of the bound ν can be guided by the discretization of the explicit operator. As an example, we compare – in Fig. 5 – the Fourier symbol of the third order finite difference $\kappa = 1/3$ advection scheme (see for example [7]) with the reduced stability domain, $\mathcal{S}^{0.5}$ for $k = 3$. The Courant number in this particular example is $\sigma = 0.35$ which is well above the value for which the scheme is expected to be SSP ($\sigma_{\text{SSP}} = 0.25$). In the table below, we will use the value $\nu = 1/3$ which corresponds fairly well to $\sigma = 0.25$ for this advection discretization.

4 Discussion and concluding remarks

In this note we have proposed and analyzed the linear stability properties of implicit-explicit methods based on the Strong Stability Preserving multistep methods introduced by Shu [11]. Of the four variant methods that we consider, one scheme stands out from the rest. The three-step scheme (16a) with $k = 3$ retains full implicit A -stability for the entire explicit stability domain. In fact, we are not aware of any other second order IMEX scheme with this property. For the other methods, we have shown $A(\alpha)$ -stability for the entire, or slightly reduced explicit stability domains.




Fig. 5. Explicit stability region $\mathcal{S}$ for the time discretization (8a), with the Fourier symbol of the third order $\kappa = 1/3$ finite difference advection scheme ($\sigma = 0.35$), and the bound ($\gamma = 0.5$) that defines the restricted stability region $\mathcal{S}^\gamma$ in Lemma 5.

It is instructive to compare with the traditional multistep IMEX methods such as the modified Crank-Nicholson/Adams-Bashforth (mCNAB) scheme and the IMEX BDF2 scheme. The mCNAB scheme introduces an algorithmic parameter, c , to the implicit integrator and can be written

$$y_{n+1} - y_n = \frac{1}{2}\Delta t \left(3f_n - f_{n-1} + (1+c)g_{n+1} + (1-2c)g_n + cg_{n-1} \right).$$

Note that $c = 0$ recovers the original Crank-Nicholson scheme for the implicit integrator, whereas other choices of c may improve the stability properties of the method, see [2,3] for a discussion. The IMEX BDF2 scheme is given by

$$3y_{n+1} - 4y_n + y_{n-1} = 2\Delta t \left(2f_n - f_{n-1} + g_{n+1} \right).$$

Both these schemes are restricted to implicit $A(\alpha)$ -stability if we require full explicit stability [3]. We summarize the implicit stability properties for all these methods in Table 1. Furthermore, all the methods appear to have comparable non-linear stability properties. Whereas the schemes proposed in this work are based on strong stability preserving methods, both the Adams-Bashforth method and the extrapolated BDF2 method was recently shown to have good monotonicity-preserving properties provided suitable starting procedures were used [6,8]. Finally, it is difficult to assess the computational cost of each method, as this may depend strongly on details of the implementation. It is however reasonable to assume that all the methods considered in

Table 1

Comparison of the implicit stability domains with full explicit stability for the schemes considered in this work and the traditional IMEX BDF2 and mCNAB methods. The results for the latter methods are taken from Reference [3].

Scheme	α
(16a) ($k = 3$)	$\pi/2$
(16a) ($k = 4$)	0.23π
(16b) ($\beta = 0, \nu = 1/3, k = 3$)	$\pi/4$
(16b) ($\beta = 0, \nu = 1/3, k = 4$)	0.15π
IMEX BDF2	0.31π
mCNAB ($c = 0$)	0
mCNAB ($c = 1/8$)	0.12π
mCNAB ($c = 1/2$)	0.23π

this study as well as the mCNAB and IMEX BDF2 schemes have comparable computational cost. The memory requirements may however be larger for the SSP-based schemes as data from more time levels need to be stored.

We will also briefly discuss IMEX Runge-Kutta methods. Ascher et al. [1] suggested that a two-stage, second-order IMEX Runge-Kutta scheme based on the trapezoidal rule could be an alternative to the CNAB method in ‘most situations’. There are also schemes in this family with very good SSP properties. Analysis of IMEX RK2 schemes shows unfortunately that there are severe restrictions on the explicit stability domain, in particular for implicit eigenvalues with large magnitude [9]. Higher order IMEX RK methods have been proposed and analyzed by Ascher et al. [1], Pareschi and Russo [10], and Higueras [5], but these have not been considered in the present work.

References

- [1] U. M. Ascher, S. J. Ruuth, and R. J. Spiteri. Implicit-explicit Runge-Kutta methods for time-dependent partial differential equations. *Appl. Numer. Math.*, 25:151–167, 1997.
- [2] U. M. Ascher, S. J. Ruuth, and B. T. R. Wetton. Implicit-explicit methods for time-dependent partial differential equations. *SIAM J. Numer. Anal.*, 32:797–823, 1995.
- [3] J. Frank, W. Hundsdorfer, and J. G. Verwer. On the stability of implicit-explicit linear multistep methods. *Appl. Numer. Math.*, 25:193–205, 1997.

- [4] S. Gottlieb, C. W. Shu, and E. Tadmor. Strong stability-preserving high-order time discretization methods. *SIAM Rev.*, 43:89–112, 2001.
- [5] I. Higueras. Strong stability for additive Runge-Kutta methods. In *Book of abstracts. ICOSAHOM 2004*. Brown University, 2004.
- [6] W. Hundsdorfer and J. Jaffré. Implicit-explicit time stepping with spatial discontinuous finite elements. *Appl. Numer. Math.*, 45:231–154, 2003.
- [7] W. Hundsdorfer, B. Koren, M. van Loon, and J. G. Verwer. A positive finite-difference advection scheme. *J. Comput. Phys.*, 117:35–46, 1995.
- [8] W. Hundsdorfer, S. J. Ruuth, and R. J. Spiteri. Monotonicity-preserving linear multistep methods. *SIAM J. Num. Anal.*, 41:605–623, 2003.
- [9] W. Hundsdorfer and J. G. Verwee. *Numerical Solution of Time-Dependent Advection-Diffusion-Reaction Equations*. Springer, 2003.
- [10] L. Pareschi and G. Russo. Implicit-explicit Runge-Kutta schemes for stiff systems of differential equations. In L. Brugnano and D. Trigiante, editors, *Recent Trends in Numerical Analysis*, volume 3 of *Advances in Computation: Theory and Practice*, pages 269–289. Nova Science, 2000.
- [11] C. W. Shu. Total-variation-diminishing time discretizations. *SIAM J. Sci. Stat. Comp.*, 9:1073–1084, 1988.